



Stony Brook Technology Center . 3400 Nesconset Highway (8 Technology Drive) . Suite 101 . Setauket, NY 11733
6144 Route 25A . Suite D21 . Wading River, NY 11792 Phone: 631-751-8700 . Fax: 631-751-5971 . www.lidigestive.com

WILLIAM J. COHN, M.D.
GARY R. BERNSTEIN, M.D.
EUGENE A. COMAN, M.D.
HOWARD A. KESCHNER, M.D.

ROBERT J. BARANOWSKI, M.D.
SETH E. PERSKY, M.D.
EMILY GLAZER, M.D.
EUGENE LICHT, M.D.

MARGARET A. CARIOLA, C.A.N.P.
KRISTINA L. DEMARCO, C.A.N.P.
HELENA C. JOSEPH, ANP-C
JILLIAN FRATEPIETRO, ANP-C

PROCEDURE DATE: _____ LI DIGESTIVE ENDOSCOPY CENTER
_____ MATHER _____ ST CHARLES

(If you need to **cancel** your procedure, please let us know **as soon as possible**.)

LIDDC: PREPOPIK INSTRUCTIONS

* FOLLOW THESE INSTRUCTIONS - NOT INSTRUCTIONS IN PREP KIT

A. MORE THAN TWO WEEKS BEFORE YOUR PROCEDURE: Inform the office (631) 751-8700 IF:

1. You are diabetic.
2. You are allergic to latex.
3. You take blood thinning products (Coumadin, Plavix, Aggrenox, Brilinta, Pradaxa, Pletal, Brilinta, Ticlid, Trental, Effient, Xarelto, Eliquis).
4. You have a prosthetic heart valve, a history of endocarditis, vascular graft, prosthetic joint replacement within the last 12 months. You may need antibiotic coverage for your procedure.
5. Make sure you have a responsible party to take you home after the procedure. We may cancel the procedure if you do not have a responsible party to take you home.

B. ONE WEEK BEFORE YOUR PROCEDURE:

1. Do not take iron, iron-containing compounds, Vitamin E, fish-oil, Advil, Motrin, ibuprofen, Aleve, naprosyn. You may take Tylenol (acetaminophen).
2. If you are taking aspirin 81 mg/day (baby aspirin) you should CONTINUE this unless otherwise instructed.
3. Blood work should be done 7-14 days prior to the procedure. A pregnancy test should be done less than 7 days prior to the procedure if you are a woman of child bearing age.
4. Avoid fruits/vegetables/supplements with seeds: e.g. tomatoes, kiwi fruit, caraway, flax, poppy, sesame and corn.

C. THE DAY BEFORE YOUR PROCEDURE:

1. **DO NOT EAT SOLID FOOD.** Drink at least 8 glasses of clear liquids. These should not be red or purple. Examples include: apple juice, water, clear broth or bouillon, Gatorade, Snapple, carbonated soda, jello, ice popsicles, black coffee, black tea (sugar or sweetener is allowed – but NO milk/creamer). It is very important that you stay well hydrated. Call the office if you have problems or questions with the prep.

-OVER-

2. **Prepopik** instructions:

A complete preparation requires 2 packets of Prepopik for oral solution taken separately, each followed by additional fluids (dosing cup provided). The 2 packets are the same and have the same contents.

a) Prepare the first 5 ounces of solution using a packet of Prepopik and drink this at 6 PM:

- Tear open the packet and dissolve the powder contents in 5 ounces (lower line on the dosing cup) of cold water
- **Stir 3 minutes – MUST BE COMPLETELY DISSOLVED**
- Drink all of the solution

b) After drinking the five ounces of Prepopik solution, drink an additional **five** 8-ounce (upper line of the dosing cup) cups of GATORADE, G-2, OR PEDIALYTE at a rate that is comfortable for you.

c) If your procedure is scheduled before 10 AM drink the second 5 ounces of solution using the other packet of Prepopik at 10 PM as you did in Step a) above.

d) If your procedure is scheduled after 10AM drink the second 5 ounces of solution using the other packet of Prepopik 6 hrs before your procedure time as you did in Step a) above.

e) After drinking the second 5 ounces of Prepopik solution, drink an additional **three** 8-ounce (upper line on the dosing cup) cups of GATORADE, G-2, OR PEDIALYTE at a rate that is comfortable for you.

3. If your procedure is scheduled in the **hospital, call the evening before** for your arrival time.

- Mather Hospital 476-2717
- St Charles Hospital 474-6441

D. THE DAY OF YOUR PROCEDURE:

1. **Do not eat solid food on the day of your procedure.** Do not drink anything in the 6 hours before your procedure. No gum or candy. You may brush your teeth. Take your usual morning medications with a sip of water.
2. Wear loose fitting clothes. Do not wear contact lenses, jewelry or make-up. Remove tongue and nose piercings.
3. Bring your inhalers and glasses case with you.
4. Contact the office if you have questions or your physical condition changes (cold, fever, illness).
5. Your responsible party will need to sign your discharge papers. You will receive discharge instructions from the nurse. You may resume your normal diet. You may not drive for the rest of the day/night. You may resume normal activities the next day.